

介護予防？－何を、どう予防するんだ？－ 参考文献一覧

【P6.老化モデル】

Busse EW: Duke University Longitudinal Studies of Aging. Z Gerontol. 26 (3) :123-8, 1993

Aging, natural death, and the compression of morbidity. Fries JF. N Engl J Med. 1980 Jul 17;303(3):130-5.

新井武志, 大淵修一・他: 地域在住高齢者の身体機能と高齢者筋力向上トレーニングによる身体機能改善効果との関係. 日老医誌 43:781-788,2006.

【P8.活動能力低下】

日本理学療法士協会:平成 20 年度老人保健事業推進費等補助金事業 介護予防事業における運動器の機能向上と生活空間等に関する調査研究事業報告書「介護予防を有効に評価するための E-SAS (無作為化比較試験デザインによる多角的検証)」2009.

【P14.痛み】

(「OARSI によるエビデンスに基づくエクスパートコンセンサスガイドライン」;2011)

Uthman OA: BMJ.2013;347:f5555 doi: 10.1136/bmj.f5555.

【P16.転倒、骨折予防】

Gillespie LD et al : Reviews 2003 , CD000340

Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson LM, Lamb SE. Interventions for preventing falls in older people living in the community. Cochrane Database Syst Rev. 2012 Sep 12;9:CD007146.

Sherrington C, Tiedemann A, Fairhall N, Close JC, Lord SR. N S W Public Health Bull. Exercise to prevent falls in older adults: an updated meta-analysis and best practice recommendations. 2011 Jun;22(3-4):78-83.

Bassey EJ, Ramsdale SJ. Increase in femoral bone density in young women following high-impact exercise. Osteoporos Int. 1994 Mar;4(2):72-5.

Bassey EJ and Ramsdale SJ : weight-bearing exercise and ground reaction forces: a 12-month randomized controlled trial of effects on bone mineral density in healthy postmenopausal women. Bone 16:469-476, 1995

Obuchi S et al: analysis of compression force in the hip joint during impulsive exercises: a preliminary study for developing exercise protocol for osteoporosis. J Jpn Physic Ther Ass 3:7-11, 2000

【P18.認知機能低下】

Suzuki T et al.: BMC Neurol. 2012, 12 : 128

Colcombe SJ, Erickson KI, Scalf PE, Kim JS, Prakash R, McAuley E, Elavsky S, Marquez DX, Hu L, Kramer AF. Aerobic exercise training increases brain volume in aging humans. J Gerontol A Biol Sci Med Sci. 2006 Nov;61(11):1166-70.

Erickson KI, Voss MW, Prakash RS, Basak C, Szabo A, Chaddock L, Kim JS, Heo S, Alves H, White SM, Wojcicki TR, Mailey E, Vieira VJ, Martin SA, Pence BD, Woods JA, McAuley E, Kramer AF. Exercise training increases size of hippocampus and improves memory. Proc Natl Acad Sci U S A. 2011 Feb 15;108(7):3017-22.

Larson EB, Wang L, Bowen JD, McCormick WC, Teri L, Crane P, Kukull W. Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. Ann Intern Med. 2006 Jan 17;144(2):73-81.

J. Carson Smith , Kristy A. Nielson , John L. Woodard , Michael Seidenberg, Stephen M. Rao: Physical Activity and Brain Function in Older Adults at Increased Risk for Alzheimer's Disease. Brain Sci. 2013, 3, 54-83

Montero-Odasso M, Bergman H, Phillips NA, Wong CH, Sourial N, Chertkow H. Dual-tasking and gait in people with mild cognitive impairment. The effect of working memory. BMC Geriatr. 2009 Sep 1;9:41.

【P20.嚥下】

藤島一郎：脳卒中の摂食・嚥下障害 第2版（医歯薬出版株式会社）

吉田剛，内山靖：脳血管障害による嚥下運動障害者の嚥下障害重症度変化と嚥下運動指標および頸部・体幹機能との関連性. 日老医誌: 43 (6), 755-760, 2006

【P22.尿失禁】

Dumoulin C, Hay-Smith J. Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women : Cochrane Database Syst Rev.2006 Jan 25;(1):CD005654.

Ueda T et al. : Urinary incontinence among community-dwelling people aged 40 years or older in Japan:prevalence, risk factors, knowledge and selfperception. Int.J.Urol . 7, 2000, 95-103

【P24.仲間との活動】

向山晶貴、中村恭子：民間フィットネスクラブ会員における運動習慣継続要因の検討. 順天堂スポーツ健康科学研究 3(3)：176-183, 2012.

西田保、渡辺俊彦、他：中高年者の運動への動機づけを促進及び阻害する要因に関する研究. デサントスポーツ科学 21：15-26, 2000